

Vegetativni prirast udomaćenih sorata masline u kolekciji u Vodnjanu 2016. godine

Krpina, Tvrtko

Master's thesis / Diplomski rad

2016

Degree Grantor / Ustanova koja je dodijelila akademski / stručni stupanj: **University of Zagreb, Faculty of Agriculture / Sveučilište u Zagrebu, Agronomski fakultet**

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:204:116652>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-04-01**



Repository / Repozitorij:

[Repository Faculty of Agriculture University of Zagreb](#)



SVEUČILIŠTE U ZAGREBU
AGRONOMSKI FAKULTET

Tvrtko Krpina

**Vegetativni prirast udomaćenih sorata
masline u kolekciji u Vodnjanu 2016.
godine**

DIPLOMSKI RAD

Zagreb, 2016

SVEUČILIŠTE U ZAGREBU
AGRONOMSKI FAKULTET

Tvrtko Krpina

**Vegetativni prirast udomaćenih sorata
masline u kolekciji u Vodnjanu 2016.
godine**

DIPLOMSKI RAD

Mentor: prof.dr.sc. Đani Benčić

Zagreb, 2016

Ovaj diplomski rad je ocijenjen i obranjen dana _____

s ocjenom _____ pred Povjerenstvom u sastavu:

1. Izv.prof.dr.sc. Đani Benčić _____

2. Prof. dr.sc. Jerko Gunjača _____

3. Izv.prof.dr.sc. Boris Duralija _____

SADRŽAJ

1. UVOD	1
1.1. Pregled literature.....	1
1.2. Osnovni podaci o predmetu istraživanja.....	2
2. CILJ ISTRAŽIVANJA	3
3. MATERIJAL I METODE ISTRAŽIVANJA	4
3.1. Nacionalna kolekcija i matični nasad u Vodnjanu.....	4
3.2. Lokacija.....	5
3.3. Klima.....	5
3.3.1. Oborine.....	6
3.3.2. Temperatura.....	6
3.4. Pedologija tla.....	7
3.5. Analiza istraživanja.....	7
5. REZULTATI I RASPRAVA	9
5.1. Ukupni promjer debla.....	9
5.2. Ukupan broj grana po stablu.....	10
5.3. Ukupna duljina grana po stablu.....	11
5.4. Usporedba rezultata s predhodnom godinom.....	12
5.5. Usporedba tabli prema promjeru debla.....	12
6. ZAKLJUČAK	13
7. LITERATURA	14

SAŽETAK

Početak travnja 2016. provedeno je istraživanje vegetativnog prirasta maslina na lokaciji Nacionalnog kolekcijuskog nasada domaćih i udomaćenih sorata maslina u Vodnjanu, čiju je sadnju 2011. godine potaknuo Agronomski fakultet u Zagrebu. Istraživanje je obuhvatilo mjerenje promjera debla, duljine grana te određivanje broja grana maslina. Istraživanje je provedeno da bi se ustvrdila vegetativna svojstva više od pedeset različitih genotipa maslina koje se već nalaze u proizvodnji u RH ili se proizvođačima nude kao nove kreacije. Među istraživanim sortama utvrđene su očekivane razlike vegetativnog prirasta. Najveći prosječni promjer debla zabilježen je kod sorte broj 7; 5,9 cm, dok je najmanji prosječni promjer debla izmjeren je među sortom masline broj 53; 3 cm. Prosječni promjer debla iznosio je 4,3 cm. Najveća prosječna du-ljina grane zabilježena je također kod sorte broj 7; 2.246 cm, dok je najmanja prosječna duljina grana izmjerena je među sortom masline broj 58; 722 cm. Prosječna duljina grana iznosila je 1.276 cm. Najveći prosječni broj grana zabilježen je kod sorte broj 7; 231, dok je najmanji prosječni broj grana izbrojen je među sortom masline broj 31; 96 grana. Prosječno su stabla u istraživanju imala 164 grane.

Ključne riječi: maslina, genotip, sorte, vegetativni prirast

ABSTRACT

The research on vegetative growth of olives, which are located at the National Collection of indigenous and domesticated varieties of olives plantation in Vodnjan, whose planting in 2011. was encouraged by Faculty of Agriculture in Zagreb, was done in the beginning of April 2016. The research included measuring trunk diameter and branches length, and also counting the number of the branches of the olive tree. Research was conducted to determine vegetative features of more than 50 different genotypes of olives that have already been producing in the Republic of Croatia or they have already been offered to manufacturers as new creations. Among the researched varieties, there was found the expected vegetative growth. The highest average trunk diameter was measured among the variety number 7; 5.9cm, while the lowest average trunk diameter was measured among the variety number 53; 3cm. The average trunk diameter of all varieties was 4.3cm. The highest average branch length was measured among the

variety number 7; 2.246cm, while the lowest average length number 58; 722cm. The average length of branches was 1,276cm. The highest number of branches was counted among the variety number 7; 231 branches, while the lowest average number of branches was counted among the variety number 31; 96 branches. The average number of branched was 164.

Keywords: olive, genotype, variety, vegetative growth